

## ONE-STEP EXTRACTION VERSUS QuEChERS FOR PESTICIDE ANALYSIS IN SELECTED FRUITS AND VEGETABLES

Darko Andjelković<sup>1</sup>, Milica Branković<sup>2\*</sup>

<sup>1</sup>University of Niš, Faculty of Agriculture, Kosancićeva 4, 37 000 Kruševac, Serbia

<sup>2</sup>University of Niš, Faculty of Science and Mathematics, Višegradska 33, 18 000 Niš, Serbia

[milica.chem@outlook.com](mailto:milica.chem@outlook.com)

This research was focused on the performance evaluation of a simple sample preparation involving acetonitrile extraction followed by liquid chromatography/mass spectrometry (LC/MS) analysis. Simplified method validation parameters, along with several other features, were compared to those of the citrate QuEChERS for 19 pesticides analyzed in four representative fruits and vegetables. The results showed comparable performances of the two methods for 5 of the 6 investigated validation parameters. The simplified method had better performance regarding the selectivity, since three analytes experienced a selectivity issue in one of four QuEChERS-treated matrices. Overall, results lead to an assumption that acetonitrile extraction could be reasonably implemented in certain cases of pesticide analysis, as an efficient and economical alternative to the official method. Since the research provides an insight into acetonitrile extraction capabilities in the domain of pesticide analysis in complex matrices, scientists, researchers or analytical practitioners can determine which method is most beneficial for a particular analysis.

**Keywords:** acetonitrile; EN 15662; HPLC; sample preparation

## ЕКСТРАКЦИЈА ВО ЕДЕН ЧЕКОР НАСПРОТИ QuEChERS ЗА АНАЛИЗА НА ПЕСТИЦИДИ ВО ИЗБРАНО ОВОШЈЕ И ЗЕЛЕНЧУК

Ова истражување беше фокусирано на евалуација на перформансите на едноставна подготовка на примерок која вклучува екстракција со ацетонитрил проследена со анализа со течна хроматографија/масена спектрометрија (LC/MS). Параметрите за валидација на поедноставениот метод, заедно со неколку други карактеристики, беа споредени со оние на цитратно пуфериран QuEChERS за 19 пестициди анализирани во четири репрезентативни видови овошје и зеленчук. Резултатите покажаа споредливи перформанси на двата метода за 5 од 6-те параметри истражени за валидација. Поедноставениот метод имаше подобри перформанси во однос на селективноста, бидејќи три аналити покажаа проблем со селективноста во една од четирите матрици третирани со QuEChERS. Општо земено, резултатите водат до претпоставка дека екстракцијата со ацетонитрил може разумно да се спроведе во одредени случаи на анализа на пестициди, како ефикасна и економична алтернатива на официјалниот метод. Бидејќи истражувањето дава увид во способностите за екстракција со ацетонитрил во доменот на анализа на пестициди во сложени матрици, научниците, истражувачите односно аналитичките практичари можат да одредат кој метод е најкорисен за одредена анализа.

**Клучни зборови:** ацетонитрил; EN 15662; HPLC; подготовка на примероци

## 1. INTRODUCTION

The diversity and complexity of fruit and vegetable contents remain the main challenge in pesticide monitoring in these matrices. Consequently, various sample preparation methods have been developed, each one with the same goal, to generate consistent high-quality results of multi-residue analysis in many plant matrices at the lowest possible pesticide concentrations. At this point, the established sample preparation method is QuEChERS. Owing to the simple steps, QuEChERS is time effective and less prone to errors, thus it represents a streamlined approach to assess multiple pesticide residues in food.

A recent literature overview shows that the established QuEChERS method and its versions are dominantly employed for pesticide analysis in fruits and vegetables.<sup>1-4</sup> The non-QuEChERS sample preparation methods,<sup>5,6</sup> simple solvent extraction methods<sup>7,8</sup> and methods comprising minimal sample preparation<sup>9-12</sup> are less frequently reviewed. A gas-liquid microextraction technique (GLME), based on the distinctive boiling points between the analytes and the interferences to achieve effective separation, has been coupled to a dispersive solid-phase extraction (*d*SPE) in a one-step sample pretreatment approach by Jin and co-workers.<sup>6</sup> The main feature of such a method is that the integrated extraction and clean-up can be performed in several minutes. This method was evaluated for 35 pesticides in apple, leek, orange and honey matrices. Guo and co-workers<sup>5</sup> implemented the reverse approach, involving carbamate pesticide extraction from tomato and apple samples onto column-packed covalent organic frameworks with acrylamide sites, eluted with acetonitrile and subjected to the liquid chromatography/ultraviolet-visible analysis (LC/UV-Vis). Solid-liquid extraction with acetonitrile was employed for the analysis of chlorpyrifos and acetamiprid in tomato peel.<sup>7,8</sup>

Methods comprising minimal sample preparation depend on the technique used for instrumental analysis. A minimal sample preparation was utilized in the screening of 6 pesticides on lemon surface by the paper-spray ionization mass spectrometry.<sup>9</sup> Pesticide detection can be achieved by rubbing the fruit surface with some tool, then exposing it to the mass spectrometer inlet. Another technique involving minimal sample preparation is the surface-enhanced Raman spectroscopy (SERS), an optimized version of Raman spectroscopy, that involves an application of gold- and/or silver-based nanoparticles onto the analytical surface to enhance the Raman signal. The technique was developed for the analysis of thiram in apples,<sup>11</sup> triazo-

phos in apples and cherry tomatoes,<sup>10</sup> and chlorpyrifos on tomato and grape surfaces.<sup>12</sup>

The objective of this study was to evaluate the performance of solvent extraction as the simplest possible sample preparation method. Evaluation included 19 pesticides, acetonitrile as an extraction solvent and four commodities (cucumber, lettuce, tomato and lemon) chosen for diverse texture, pigmentation and acidity.

The performances of pesticide analysis in the selected fruits and vegetables were compared to those of the citrate-buffered QuEChERS followed by dispersive solid phase extraction (*d*SPE) with PSA. Contrary to the practice of analysis in multiple reaction monitoring (MRM) mode, analytes were detected and quantified in the full scan MS<sup>1</sup> mode. Since the goal was to evaluate and compare the performances of the two methods, the simplification in mass analysis was intentionally implemented. Furthermore, if the screening itself is a purpose, recording a sample's spectrum in full scan mode will offer a possibility for the retroactive analysis of some other analytes, not covered by the current plan of analysis.

## 2. MATERIALS AND METHODS

### 2.1. Chemicals and consumables

Formic acid (FA) (98 %) and high purity pesticide standards (acetamiprid, azoxystrobin, boscalid, buprofezin, chlorpyrifos, cyprodinil, difenoconazole, fenhexamid, imidacloprid, kresoxim-methyl, metsulfuron-methyl, propiconazole, pyraclostrobin, pyrimethanil, pyriproxyfen, tebuconazole, thiacloprid, thiamethoxam, trifloxystrobin) were produced by Sigma-Aldrich® (St. Louis, Missouri, USA). Ammonium formate (AMF) (98 %), deionized water and high-performance liquid chromatography (HPLC) grade ethanol were produced by Carlo Erba (Emmendingen, Germany). HPLC grade methanol (MeOH) and HPLC grade acetonitrile (AcN) were produced by J.T. Baker, Thermo Fisher Scientific (Waltham, Massachusetts, USA). For the sample preparation, procedure prepacked Hilleium QuEChERS extraction pouches (1 g of NaCl, 4 g of MgSO<sub>4</sub>, 1 g of trisodium citrate dihydrate and 0.5 g disodium hydrogen citrate) and Hilleium QuEChERS dispersion kits (25 mg PSA and 150 mg MgSO<sub>4</sub>) were used. Syringe microfilters (Nylon Hydrophilic 0.22 μm) were produced by Membrane Solutions (Auburn, Washington, USA).

Fresh fruits and vegetables (tomato, cucumber, lettuce, and lemon) were purchased in a local supermarket.

## 2.2. Instruments and instrumental parameters

**Appliances.** For high purity standards weighing procedures, the analytical balance Sartorius BP110S (Göttingen, Germany) was used. The sample preparation procedure used the following appliances: blender 0.9 l BL142A by TEFAL (Rumilly, Haute-Savoie, France); balance (acc.  $\pm$  0.01 g) KB 2000-2N by KERN & SOHN GmbH (Balingen, Germany); and centrifuge Jouan C4i by Thermo Fisher Scientific. To facilitate the extraction, Digital Vortex-Genie 2 by Scientific Industries (Bohemia, New York, USA) was used. Nitrogen (99 %) was supplied by a nitrogen generator by PEAK Scientific (Glasgow, Scotland, UK).

**Analytical instruments.** Instrumental analysis was performed on an LC/MS system including a Surveyor autosampler by Thermo Finnigan LLC (San Jose, California, USA), and an Accela MS pump and LTQ XL mass spectrometer with linear ion trap analyzer by Thermo Fischer Scientific. Analytes were separated on a Hypersil GOLD column (C<sub>18</sub>, 150 mm  $\times$  2.1 mm, particle size 3  $\mu$ m), ionized in ESI+ ionization mode and monitored in MS<sup>1</sup> full scan mode (scan range  $m/z$  150–600). Data was acquired and analyzed by Thermo Xcalibur™ software, version 2.1.0, SP1.1160.

**Instrumental parameters.** Ten microliters of sample were loaded on to a column in a partial loop injection mode and eluted with a mixture containing eluent A (buffer solution –0.1 % of FA and 0.03 % of AMF in water) and eluent B (MeOH), following the gradient: 0 min (90 % A), 0–2 min (90 % A), 2–7 min (30 % A), 7–30 min (30 % A), 30–35 min (90 % A) and 35–40 min (90 % A) with a flow rate equal to 300  $\mu$ l min<sup>-1</sup>. The chromatographic column was kept at thermostatic conditions at 25 °C. ESI source parameters were: sheath gas = 21 arbitrary units; auxiliary gas = 18 arb; *I* (spray voltage) = 5 kV; capillary *T* = 275 °C. Protonated molecular ions of tested analytes cover the  $m/z$  range from 200 to 409 (Table S1), therefore, the ion optics were optimized according to the pesticides' ions from the lower, middle and higher part of the  $m/z$  range, that is, according to the ions of pyrimethanil, buprofezin and trifloxystrobin. Tuned parameters meeting optimal detectability for all analytes were chosen.

## 2.3. Procedures

**Stock preparation.** Single-pesticide stock solutions (1 mg ml<sup>-1</sup> each) were prepared by dissolving high purity pesticide standards in ethanol. Multi-pesticide solutions were prepared by mixing and diluting single stocks in ethanol.

**Sample homogenization.** Lemon, tomato, cucumber and lettuce, one kilogram of each, were cut and homogenized by blending for 5 min. All parts (flesh and peel) of the tomato, lettuce and cucumber were included in analysis. For lemon samples, only the flesh was used.

**Sample preparation – extraction with AcN.** Ten grams of homogenate were extracted with 10 ml of AcN. Extracts were centrifuged (10 min/3000 rpm) and supernatants were microfiltered prior to instrumental analysis. In the case of procedural standards preparation, the homogenate portion was spiked prior to the solvent extraction.

**Sample preparation – citrate-buffered QuEChERS (EN 15662).** Ten grams of homogenate were extracted with 10 ml of AcN, after which an extraction pouch was added. The mixture was immediately vortexed for one minute and centrifuged (10 min/3000 rpm). A supernatant aliquot was subjected to a dispersive extraction by the addition of one dispersion kit per ml of supernatant. The mixture was vortexed for one minute and centrifuged (10 min/3000 rpm). Supernatant was microfiltered prior to instrumental analysis. In the case of procedural standards preparation, the homogenate portion was spiked prior to the solvent extraction.

**Validation study.** Samples obtained from the market were initially screened for tested pesticides residues. Since none of the targeted pesticides was detected, the samples were considered blank samples and used for validation procedures. The matrix effect (ME) was evaluated by comparing the slopes of the calibration curves (0.01–15.00  $\mu$ g ml<sup>-1</sup>) of solvent-based and matrix-based multi-pesticide standards. The linear range of methods was evaluated with 7 procedural standards in the concentration range 0.00–15.00 mg kg<sup>-1</sup>. Chromatographic repeatability was evaluated from the successive injections of ten procedural-based (5.00 mg kg<sup>-1</sup>) and ten solvent-based standards (5.00  $\mu$ g ml<sup>-1</sup>). Trueness and precision were evaluated at three concentration levels (0.50, 5.00 and 15.00 mg kg<sup>-1</sup>), each level at 5 replicates. Detection limits (DLs) were evaluated by the S/N criterion. The concentration of spiked sample that produced an analyte peak with  $S/N = 3$  is established as the method's DL. In cases where the spiked samples produced an  $S/N$  ratio other than 3, the DL was estimated by a proportion involving the procedural standard with S/N ratio closest to 3.

## 3. RESULTS AND DISCUSSION

A prerequisite for a successful analysis is chromatographic or mass separation of analytes. Among the targeted analytes, boscalid and propiconazole would express a selectivity issue if they

were not chromatographically separated, due to the closeness of  $MH^+$   $m/z$  values (Table S1).

Both one-step solvent extraction and the QuEChERS method expressed similar performances in terms of analyte recovery. Recoveries of the targeted analytes from tomato and lemon treated with the QuEChERS were within the limits (70–120 %) (Figs. S1 and S4). In lettuce and cucumber (Figs S2 and S3), at 0.50 mg kg<sup>-1</sup> spike level, the recovery of imidacloprid was much below 70 %. Recoveries of the targeted analytes from each matrix treated with the simplified method (AcN extraction) were within the limits (70–120 %) (Figs. S1–S4). Considering the average recovery, both methods demonstrated satisfactory performance for each analyte (Fig. 1). The average recovery of thi-

amethoxam, after AcN extraction from lemon samples was less than 70 % (Fig. 1), however, out-of-limits recovery was 68 %. According to SAN-TE/11312/2021,<sup>13</sup> recoveries out of the 70–120 % range are acceptable if they are consistent (RSD < 20 %), but the mean recovery must not be lower than 30 % or higher than 140 %.

The detection limits of the two methods were comparable, and in most cases below or equal to the pesticide maximum residue limit (MRL) (Fig. 1). In three cases, the DLs were higher than the MRLs. The DLs of both methods were slightly higher than fenhexamid MRL in lemon; the DL of the QuEChERS was higher than cyprodinil MRL in lemon (Fig. 1).




Fig. 1. Main performances overview of two methods for the targeted pesticide analysis in selected fruits and vegetables

Both methods demonstrated similar performances regarding the matrix effect and linearity range. Strong signal suppression was observed for each analyte in each tested matrix, regardless of the sample preparation procedure (Figures S1-S4). Analyte signals were suppressed by at least 37 %. Methods were linear in the tested analytical range (0.00 – 15.00 mg kg<sup>-1</sup>), with back-calculated concentration deviations < 20 % and correlation coefficients > 0.90 (Table 1).

Significant pesticide retention time shifts were observed in sample extracts, regardless of the sample preparation procedure (Figure S5). The shift ranged from 0.12 to 1.11 min. The tolerable limit of ± 0.10 min was exceeded by 8 analytes in tomato, 4 in lettuce and cucumber and 12 analytes in lemon, all treated with the QuEChERS. The exceedance rate in samples treated with the simplified method was similar i.e., the retention time shift of 3 analytes in tomato, 5 in lettuce, 12 in cucumber and 11 in lemon exceeded the limits. The strongest shift was noticed in the lemon matrix, which stands out with the lowest pH value of the final extracts.

In terms of method selectivity, lemon was identified as a difficult matrix. A selectivity issue was observed for pyraclostrobin, tebuconazole and fenhexamid in the QuEChERS-treated lemon (Figure S6) and for fenhexamid in AcN-treated lemon (Figure S7). The issue for pyraclostrobin and fenhexamid was related to the overlay of analyte chromatographic peaks with the peaks of interferences. The issue for tebuconazole was related to the overlay of its peak with the peak of buprofezin, due to the heavy retention time shift of tebuconazole in lemon. Interestingly, the selectivity issue for pyraclostrobin and tebuconazole could be exclusively observed in the QuEChERS extracts.

Analytical data for herein investigated methods and for similar methods found in the literature are generally in mutual agreement. Melton and Taylor<sup>4</sup> implemented the same QuEChERS procedure for the analysis of buprofezin and chlorpyrifos in lemon by gas chromatography/mass spectrometry (GC/MS), which resulted in more than 94 % of recovered analytes and a DL of 0.01 mg kg<sup>-1</sup>. Martinez Bueno et al.<sup>3</sup> also implemented the citrate QuEChERS for multi-pesticide LC/MS analysis in lettuce, but the *d*SPE step was performed with C<sub>18</sub>. Percentages of recovered analytes

ranged from 77 for azoxystrobin to 91 for thiacloprid. Another variation of the citrate QuEChERS followed by the *d*SPE with PSA and ENVI-Carb was implemented by Ferracane et al.<sup>1</sup> for the analysis of chlorpyrifos in tomatoes, cucumbers and lettuce by flow-modulated GC/MS. The recovery of chlorpyrifos was higher than 99 % in each matrix. The lowest DL of 1.8 µg kg<sup>-1</sup> was established for cucumbers. Mahdavi et al.<sup>2</sup> implemented the dispersive solid-phase extraction with primary-secondary amine (PSA) for pesticide LC/MS analysis in cucumbers, but as a part of the acetate-buffered QuEChERS. Ten of more than 50 investigated pesticides matched our study. The recovery ranged from 74 % for kresoxim-methyl to 107 % for propiconazole. The lowest and the highest DL of 0.002 and 0.01 mg kg<sup>-1</sup> were established for kresoxim-methyl and imidacloprid, respectively. Çatak and Tiriyaki<sup>14</sup> implemented the same acetate-buffered QuEChERS for GC/MS analysis of chlorpyrifos and acetamiprid in cucumbers. A recovery higher than 80 % and LOQs of 2 and 10 µg kg<sup>-1</sup> for acetamiprid and chlorpyrifos, respectively, were established. Dashtbozorgi et al.<sup>15</sup> applied a dispersive liquid-liquid microextraction technique for the extraction and pre-concentration of acetamiprid, imidacloprid, azoxystrobin and 17 other pesticide residues from QuEChERS extracts of tomato and cucumber. This combined procedure resulted in recovery of around 100 % and DLs ranging from 3.9 to 9.4 µg kg<sup>-1</sup>.

In addition to our study, acetonitrile-based solvent extractions were implemented by Moura et al.<sup>8</sup> and Hegazy et al.<sup>7</sup> for the analysis of chlorpyrifos and acetamiprid in tomatoes. The detection techniques, however, were the paper-spray ionization mass spectrometry (chlorpyrifos) and the LC/UV-Vis (acetamiprid), providing the DL of 0.01 ppm and 0.03 µg ml<sup>-1</sup>, respectively. In each case, pesticide recovery was higher than 94 %. A different solvent extraction procedure was implemented by Mohamed et al.<sup>16</sup> for the analysis of chlorpyrifos and five other pesticides in tomatoes and cucumbers by gas chromatography/flame ionization detection (GC/FID). Pesticides were extracted from samples in a successive extraction with acetone and dichloromethane, after which the extract was cleaned-up on a Florisil® stationary phase. Achieved DLs ranged from 0.001 (chlorpyrifos) to 0.20 mg kg<sup>-1</sup> (profenofos).

Table 1

Linear regression parameters ( $y = ax + b$ , conc. range 0.00 – 15.00 mg kg<sup>-1</sup>, 7 points) for EN 15662 and AcN method

	Method	Tomato			Cucumber			Lettuce			Lemon		
		a	b	R <sup>2</sup>	a	b	R <sup>2</sup>	a	b	R <sup>2</sup>	a	b	R <sup>2</sup>
Thiametoxam	EN 15662	49,942	607	0.999 <sub>9</sub>	39,064	4,578	0.997 <sub>1</sub>	45,668	-5,647	0.999 <sub>0</sub>	23,806	8,495	0.993 <sub>8</sub>
	AcN	29,397	-4,759	0.998 <sub>9</sub>	22,034	-1,084	0.999 <sub>8</sub>	25,274	4,890	0.984 <sub>3</sub>	21,210	7,403	0.996 <sub>4</sub>
Acetamiprid	EN 15662	486,466	248,738	0.983 <sub>4</sub>	379,027	278,482	0.981 <sub>7</sub>	425,279	587,260	0.981 <sub>5</sub>	165,015	165,89 <sub>7</sub>	0.953 <sub>6</sub>
	AcN	312,583	237,557	0.995 <sub>2</sub>	223,458	133,826	0.985 <sub>9</sub>	272,525	150,852	0.987 <sub>6</sub>	145,965	21,718	0.998 <sub>4</sub>
Thiacloprid	EN 15662	122,297	33,645	0.995 <sub>2</sub>	87,546	36,645	0.991 <sub>0</sub>	116,744	30,588	0.995 <sub>2</sub>	51,924	20,519	0.993 <sub>1</sub>
	AcN	76,465	9,992	0.999 <sub>5</sub>	49,158	16,813	0.995 <sub>8</sub>	63,079	11,697	0.995 <sub>9</sub>	29,673	11,311	0.996 <sub>9</sub>
Imidacloprid	EN 15662	21,500	29,936	0.988 <sub>1</sub>	12,056	7,919	0.990 <sub>7</sub>	14,559	18,370	0.985 <sub>0</sub>	7,042	6,694	0.999 <sub>2</sub>
	AcN	12,383	11,748	0.997 <sub>6</sub>	7,176	4,941	0.994 <sub>0</sub>	13,209	27,382	0.917 <sub>8</sub>	5,332	2,489	0.996 <sub>5</sub>
Metsulfuron methyl	EN 15662	38,308	147,477	0.911 <sub>4</sub>	22,447	99,689	0.948 <sub>2</sub>	22,974	123,848	0.907 <sub>0</sub>	15,644	63,143	0.959 <sub>5</sub>
	AcN	25,804	103,910	0.979 <sub>7</sub>	18,556	70,425	0.955 <sub>6</sub>	17,295	175,560	0.932 <sub>0</sub>	9,075	83,970	0.995 <sub>8</sub>
Pyrimethanil	EN 15662	1,170,5 <sub>66</sub>	926,818	0.990 <sub>1</sub>	742,540	756,042	0.991 <sub>1</sub>	993,453	223,656	0.999 <sub>6</sub>	403,327	23,229	0.999 <sub>9</sub>
	AcN	667,633	73,335	0.999 <sub>8</sub>	353,311	73,690	0.999 <sub>2</sub>	527,774	136,482	0.998 <sub>9</sub>	273,318	38,903	0.999 <sub>8</sub>
Azoxystrobin	EN 15662	1,576,2 <sub>30</sub>	1,241,2 <sub>36</sub>	0.966 <sub>0</sub>	982,868	754,357	0.969 <sub>7</sub>	1,303,2 <sub>63</sub>	2,373,8 <sub>00</sub>	0.981 <sub>0</sub>	624,781	509,64 <sub>0</sub>	0.975 <sub>3</sub>
	AcN	1,047,7 <sub>86</sub>	641,847	0.985 <sub>0</sub>	642,729	403,148	0.976 <sub>5</sub>	926,415	668,696	0.976 <sub>3</sub>	596,502	380,28 <sub>6</sub>	0.977 <sub>1</sub>
Cyprodinil	EN 15662	2,944,7 <sub>60</sub>	1,610,9 <sub>05</sub>	0.991 <sub>4</sub>	1,826,1 <sub>60</sub>	1,099,1 <sub>91</sub>	0.990 <sub>7</sub>	2,753,0 <sub>17</sub>	1,483,3 <sub>88</sub>	0.991 <sub>9</sub>	1,347,7 <sub>05</sub>	578,01 <sub>7</sub>	0.989 <sub>2</sub>
	AcN	1,912,9 <sub>74</sub>	1,081,9 <sub>59</sub>	0.992 <sub>8</sub>	1,101,3 <sub>83</sub>	945,764	0.992 <sub>3</sub>	1,601,0 <sub>24</sub>	1,045,9 <sub>06</sub>	0.988 <sub>5</sub>	857,576	927,46 <sub>8</sub>	0.980 <sub>1</sub>
Boscalid	EN 15662	164,078	568,662	0.949 <sub>2</sub>	105,761	509,623	0.944 <sub>9</sub>	158,072	417,802	0.959 <sub>2</sub>	73,347	301,37 <sub>0</sub>	0.979 <sub>9</sub>
	AcN	102,244	526,804	0.951 <sub>6</sub>	47,261	672,624	0.969 <sub>1</sub>	74,009	595,158	0.976 <sub>3</sub>	37,466	584,88 <sub>4</sub>	0.966 <sub>2</sub>
Fenhexamid	EN 15662	332,842	55,619	0.998 <sub>4</sub>	200,520	29,784	0.998 <sub>9</sub>	288,272	27,730	0.999 <sub>7</sub>	61,730	8,002	1.000 <sub>0</sub>
	AcN	201,886	-2,966	0.999 <sub>5</sub>	108,899	12,473	0.999 <sub>3</sub>	157,094	31,708	0.998 <sub>0</sub>	106,134	16,633	0.999 <sub>6</sub>
Kresoxim methyl	EN 15662	115,839	26,132	0.997 <sub>5</sub>	81,169	7,792	0.997 <sub>4</sub>	112,297	-35,606	0.999 <sub>2</sub>	55,463	3,860	0.999 <sub>7</sub>
	AcN	67,531	-9,810	0.999 <sub>2</sub>	44,782	-5,484	0.999 <sub>2</sub>	55,713	-1,362	0.996 <sub>7</sub>	44,333	27,402	0.996 <sub>5</sub>
Tebuconazole	EN 15662	456,056	44,800	0.999 <sub>5</sub>	335,944	46,944	0.999 <sub>5</sub>	405,717	15,865	0.999 <sub>9</sub>	49,270	4,971	0.999 <sub>6</sub>
	AcN	264,252	-15,632	0.999 <sub>3</sub>	180,366	4,045	0.999 <sub>9</sub>	212,257	42,837	0.998 <sub>1</sub>	161,520	349	0.999 <sub>9</sub>
Propiconazole	EN 15662	506,139	55,230	0.999 <sub>5</sub>	359,105	39,303	0.999 <sub>6</sub>	462,719	25,161	0.999 <sub>9</sub>	249,045	24,293	0.999 <sub>7</sub>
	AcN	298,162	-2,832	0.999 <sub>7</sub>	198,263	9,269	0.999 <sub>9</sub>	241,819	108,716	0.993 <sub>0</sub>	178,305	68,814	0.995 <sub>5</sub>
Pyraclostrobin	EN 15662	386,406	200,652	0.989 <sub>5</sub>	240,437	81,509	0.995 <sub>2</sub>	353,274	170,541	0.997 <sub>2</sub>	112,019	53,701	0.992 <sub>6</sub>
	AcN	230,295	45,069	0.997 <sub>8</sub>	140,144	32,557	0.997 <sub>8</sub>	183,003	105,754	0.995 <sub>1</sub>	92,956	61,009	0.978 <sub>6</sub>
Buprofezin	EN 15662	2,104,0 <sub>04</sub>	1,128,6 <sub>57</sub>	0.982 <sub>6</sub>	1,408,6 <sub>29</sub>	621,707	0.989 <sub>3</sub>	2,017,3 <sub>41</sub>	702,916	0.992 <sub>1</sub>	881,980	275,50 <sub>3</sub>	0.994 <sub>9</sub>
	AcN	1,295,5 <sub>27</sub>	361,312	0.996 <sub>6</sub>	860,423	232,692	0.994 <sub>9</sub>	1,186,9 <sub>21</sub>	410,927	0.991 <sub>2</sub>	705,379	256,71 <sub>5</sub>	0.991 <sub>4</sub>
Difenoconazole	EN 15662	706,205	-43,422	0.998 <sub>2</sub>	504,815	-75,625	0.999 <sub>2</sub>	664,133	-	0.999 <sub>3</sub>	329,484	9,350	1.000 <sub>0</sub>
	AcN	408,421	-	0.998 <sub>6</sub>	266,223	-	0.999 <sub>9</sub>	323,682	282,868	0.952 <sub>2</sub>	213,260	-	0.998 <sub>8</sub>
Tri-floxystrobin	EN 15662	1,675,4 <sub>20</sub>	366,819	0.997 <sub>3</sub>	1,132,7 <sub>14</sub>	311,321	0.998 <sub>7</sub>	1,584,1 <sub>88</sub>	265,710	0.999 <sub>3</sub>	802,219	166,18 <sub>7</sub>	0.999 <sub>9</sub>

		Tomato			Cucumber			Lettuce			Lemon		
Pyriproxyfen	AcN	1,015,2 59	65,089	0.999 9	590,471	255,918	0.998 7	822,058	314,827	0.996 7	541,449	159,25 8	0.996 8
	EN 15662	1,716,3 52	539,341	0.993 7	1,117,8 34	303,442	0.997 3	1,574,5 90	351,082	0.998 6	767,966	110,02 8	0.999 7
Chorpyrifos	AcN	1,019,9 48	69,890	0.999 8	616,235	106,065	0.998 1	852,834	232,179	0.995 3	546,597	134,72 3	0.996 2
	EN 15662	370,699	13,928	0.996 7	234,163	34,077	0.998 9	310,092	91,233	0.994 1	178,660	-1,059	0.999 9
	AcN	182,143	11,644	0.993 3	114,122	32,707	0.994 5	142,877	63,797	0.989 1	114,403	17,945	0.998 4

#### 4. CONCLUSION

Performances of the first step of the QuEChERS sample preparation procedure, which is the acetonitrile extraction, were evaluated for 19 pesticides in four representatives of fruits and vegetables and were compared to the performances of the citrate-buffered QuEChERS. Both methods expressed comparable performances regarding most of the validated parameters, including the matrix effect, chromatographic repeatability, recovery, detection limits and linearity. Better performances regarding the selectivity were expressed in the acetonitrile extraction method, since 3 analytes experienced a selectivity issue in QuEChERS-treated lemon. In the end, the comparable performances of the simplified method and QuEChERS reasonably qualify the simplified method for implementation in certain cases of pesticide analysis, as an efficient and economical alternative to the official method.

**Acknowledgements.** This work was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (contract no. 451-03-47/2023-01/200124 and 451-03-47/2023-01/200383).

#### REFERENCES

- Ferracane, A.; Zoccali, M.; Cacciola, F.; Grazia Salerno, T. M.; Tranchida, P. Q.; Mondello, L., Determination of multi-pesticide residues in vegetable products using a "reduced-scale" QuEChERS method and flow-modulated comprehensive two-dimensional gas chromatography-triple quadrupole mass spectrometry. *J. Chrom. A*, **2021**, *1645*:462126. <https://doi.org/10.1016/j.chroma.2021.462126>
- Mahdavi, V.; Eslami, Z.; Gordan, H.; Ramezani, S.; Peivasteh-roudsari, L.; Ma'mani, L.; Khaneghah, A. M., Pesticide residues in green-house cucumber, cantaloupe, and melon samples from Iran: A risk assessment by Monte Carlo Simulation. *Environ. Res.* **2022**, *206*:112563. <https://doi.org/10.1016/j.envres.2021.112563>
- Martinez Bueno, M. J.; Valverde, M. G.; del Mar Gomez-Ramos, M.; Valverde, A.; Martinez Galera, M.; Rodriguez Fernandez-Alba, A., Monitoring of pesticide residues in crops irrigated with reclaimed water by a multiresidue method based on modified QuEChERS. *Anal. Methods* **2021**, *13* (36):4131–4142. <https://doi.org/10.1039/D1AY00845E>
- Melton, L. M.; Taylor, M. J., Use of a deactivated PTV injector liner and GCMS/MS for the quantitative determination of multiple pesticide residues in fruit and vegetables. *MethodsX* **2021**, *8*:101180. <https://doi.org/10.1016/j.mex.2020.101180>
- Guo, H.; Chen, A.; Zhou, J.; Li, Y.; He, X.; Chen, L.; Zhang, Y., Efficient extraction and determination of carbamate pesticides in vegetables based on a covalent organic frameworks with acylamide sites. *J. Chrom. A* **2022**, *1664*:462799. <https://doi.org/10.1016/j.chroma.2021.462799>
- Jin, X.; Kaw, H. Y.; Liu, Y.; Zhao, J.; Piao, X.; Jin, D.; He, M.; Yan, X.P.; Zhou, J.; Li, D., One-step integrated sample pretreatment technique by gas-liquid microextraction (GLME) to determine multi-class pesticide residues in plant-derived foods. *Food Chem.* **2022**, *367*: 130774. <https://doi.org/10.1016/j.foodchem.2021.130774>
- Hegazy, A. M.; Abdelfatah, R. M.; Mahmoud, H. M.; Elsayed, M. A., Development and validation of two robust simple chromatographic methods for estimation of tomatoes specific pesticides' residues for safety monitoring prior to food processing line and evaluation of local samples. *Food Chem.* **2020**, *306*:125640. <https://doi.org/10.1016/j.foodchem.2019.125640>
- Martins Moura, A. C.; Neves Lagoa, I.; Fernandes Cardoso, C.; Dos Reis Nascimento, A.; Pereira, I.; Gontijo Vaz, B., Rapid monitoring of pesticides in tomatoes (*Solanum lycopersicum* L.) during pre-harvest intervals by paper spray ionization mass spectrometry." *Food Chem.* **2020**, *310*: 125928. <https://doi.org/10.1016/j.foodchem.2019.125938>
- Chen, K.H.; Li, Y.C.; Sheu F.; Lin, C. H., Rapid screening and determination of pesticides on lemon surfaces using the paper-spray mass spectrometry integrated via thermal desorption probe. *Food Chem.* **2021**, *363*: 130305. <https://doi.org/10.1016/j.foodchem.2021.130305>
- Gong, X.; Tang, M.; Gong, Z.; Qiu, Z.; Wang, D.; Fan, M., Screening pesticide residues on fruit peels using portable Raman spectrometer combined with adhesive tape sampling. *Food Chem.* **2019**, *295*:254–258. <https://doi.org/10.1016/j.foodchem.2019.05.127>
- Jiao, A.; Dong, X.; Zhang, H.; Xu, L.; Tian, Y.; Liu, X.; Chen, M., Construction of pure worm-like AuAg nano-chains for ultrasensitive SERS detection of pesticide residues on apple surfaces. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **2019**, *209*:241–247. <https://doi.org/10.1016/j.saa.2018.10.051>
- Subramaniam, T.; Kesavan, G., Coherently designed sustainable SERS active substrate of Ag/TiO<sub>2</sub> hybrid nanostructures for excellent ultrasensitive detection of

- chlorpyrifos pesticide on the surface of grapes and tomatoes. *J. Food Comp. Anal.* **2022**, *106*:104330. <https://doi.org/10.1016/j.jfca.2021.104330>
- (13) Analytical quality control and method validation procedures for pesticide residues analysis in food and feed: SANTE/11312/2021. [https://www.eurlpesticides.eu/userfiles/file/EurlALL/SANTE\\_11312\\_2021.pdf](https://www.eurlpesticides.eu/userfiles/file/EurlALL/SANTE_11312_2021.pdf) (accessed 2023-08-25)
- (14) Çatak, H.; Tiryaki, O., Insecticide residue analyses in cucumbers sampled from Çanakkale open markets. *Türk. entomol. derg.* **2020**, *44*: 449–460. <http://dx.doi.org/10.16970/entoted.767482>
- (15) Dashtbozorgi, Z.; Kazem Ramezanib, M.; Waqif-Husaina, S., Optimization and validation of a new pesticide residue method for cucumber and tomato using acetonitrile-based extraction-dispersive liquid-liquid microextraction followed by liquid chromatography-tandem mass spectrometry. *Anal. Methods* **2013**, *5*: 1192–1198. <https://doi.org/10.1039/C2AY26287H>
- (16) Mohamed, A. O.; Mater, A. A.; Hammad, A. M. A.; Ishag, A. E. S. A.; El Tayeb, E. M.; Dahab, A. A., Pesticide residues detected on tomato and cucumber fruits grown in greenhouse farms in Khartoum State, Sudan. *Int. J. Life Sci. Res.* **2018**, *6*: 472–481.