

## MECHANOCHEMICAL SYNTHESIS AND SOLID-STATE CHARACTERIZATION OF MOLECULAR SALTS OF PYRIDOXINE: VIBRATIONAL SPECTROSCOPIC AND THERMAL CONSIDERATION\*

Aleksandar Cvetkovski<sup>1\*</sup>, Ljupcho Pejov<sup>2,3,4</sup>, Monika Stojanovska Pecova<sup>5</sup>,  
Gjorgji Petrushevski<sup>6</sup>, Petre Makreski<sup>2</sup>

<sup>1</sup> Faculty of Medical Sciences, Goce Delcev University, 2000 Štip, North Macedonia

<sup>2</sup> Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University in Skopje, Skopje, North Macedonia

<sup>3</sup> Department of Chemistry, Bioscience and Environmental Engineering,


Faculty of Science and Technology, University of Stavanger, 4021 Stavanger, Norway

<sup>4</sup> The Polytechnic School, Ira A. Fulton Schools of Engineering, Arizona State University, AZ, USA

<sup>5</sup> Research and Development, Alkaloid AD, Blvd. Aleksandar Makedonski 12, Skopje, North Macedonia


<sup>6</sup> Quality Control Department, Alkaloid AD, Blvd. Aleksandar Makedonski 12, Skopje, North Macedonia

[aleksandar.cvetkovski@ugd.edu.mk](mailto:aleksandar.cvetkovski@ugd.edu.mk)



\* Dedicated on the occasion of the Golden Jubilee of the *Macedonian Journal of Chemistry and Chemical Engineering*

**Fig. 1S.** Calculated XRPD patterns of PN/SA salt (determined structure by single-crystal XRD) compared to the experimental XRPD patterns of sample from solid mixture **1** (PN/SA in 1:1 M/M), PN base and SA).



**Fig. 2S.** Calculated XRPD patterns of PN/FA·H<sub>2</sub>O salt (determined structure by single-crystal XRD) compared to the experimental XRPD patterns of sample from solid mixture **2** (PN/FA in 1:1 M/M), PN base and FA.