

MECHANOCHEMICAL SYNTHESIS AND SOLID-STATE CHARACTERIZATION OF MOLECULAR SALTS OF PYRIDOXINE: VIBRATIONAL SPECTROSCOPIC AND THERMAL CONSIDERATION*

Aleksandar Cvetkovski^{1*}, Ljupcho Pejov^{2,3,4}, Monika Stojanovska Pecova⁵,
Gjorgji Petrushevski⁶, Petre Makreski²

¹ Faculty of Medical Sciences, Goce Delchev University, 2000 Štip, North Macedonia

² Institute of Chemistry, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University in Skopje, Skopje, North Macedonia


³ Department of Chemistry, Bioscience and Environmental Engineering, Faculty of Science and Technology, University of Stavanger, 4021 Stavanger, Norway

⁴ The Polytechnic School, Ira A. Fulton Schools of Engineering, Arizona State University, AZ, USA

⁵ Research and Development, Alkaloid AD, Blvd. Aleksandar Makedonski 12, Skopje, North Macedonia

⁶ Quality Control Department, Alkaloid AD, Blvd. Aleksandar Makedonski 12, Skopje, North Macedonia

aleksandar.cvetkovski@ugd.edu.mk



* Dedicated on the occasion of the Golden Jubilee of the *Macedonian Journal of Chemistry and Chemical Engineering*

Fig. 1S. Calculated XRPD patterns of PN/SA salt (determined structure by single-crystal XRD) compared to the experimental XRPD patterns of sample from solid mixture 1 (PN/SA in 1:1 M/M), PN base and SA).




Fig. 2S. Calculated XRPD patterns of PN/FA·H₂O salt (determined structure by single-crystal XRD) compared to the experimental XRPD patterns of sample from solid mixture 2 (PN/FA in 1:1 M/M), PN base and FA.